



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Experimentelle Untersuchungen über die Helligkeit der Farben.
 EDUARD GRUBER. Wundt's Philos. Studien., IX. 1893, 429-446.

The method used by Gruber for determining the relative brightness of colors was the natural one of direct comparison, applied, however, in the form of the method of minimal change. Two color-tops, one carrying a gray disk, *i. e.*, a black and white one (*A*), and the other a colored disk (*B*), were set up before a dark background. The gray disk (*A*) was at first set decidedly too dark and gradually brightened till it seemed to the observer of equal brightness with the colored disk (*B*). The condition of *A* was noted and then made much too light and gradually darkened till a match was again reached. The average of the two determinations, repeated several times on each side, gave the brightness of the color. The experiments made in this way, after a little practice, yielded excellent results, except in a few cases where the observers seem to have become habituated to a particular intensity of gray and to have judged by its return rather than by an unbiased comparison. For the special precautions employed the original must be consulted.

The author also investigated the Purkinje phenomenon and the effect of changes in saturation on brightness. The experiments on the first, so far as they went, gave results in accord with the similar experiments of Hillebrand. With decreased illumination, the blue and green not only lost less in brightness, as compared with the red and yellow, but less even than the neutral gray. The effect of change in saturation was tested by replacing a portion of the color on *A* by an equally luminous gray, but no change in the total brightness was to be observed. Experiments on a color-blind observer (red-green blind), made in the hope of deciding whether green appeared to him relatively brighter than it did to an observer with a normal eye, which might be expected on Hering's theory, unfortunately had to be discontinued before clear indications appeared. It is, perhaps, only fair to add that the same method of measuring the brightness of colors was used by Rood more than fifteen years ago; see *American Journal of Science*, Ser. 3, XV. 1878, 81-82.
 E. C. S.

On a Color System. O. N ROOD. *American Journal of Science*, Ser. 3, XLIV. 1892, 263-270.

In this article Professor Rood describes a method of working out a reproducible color system with the color-top, provided that there is at hand a single disk of known hue and power of saturation, *i. e.*, efficiency in the formation of gray when mixed with its complement. By combination with the standard disk, the power of saturation of its complement and of colors differing but slightly from that, are determined, and from these in turn the powers of saturation of other disks, till a considerable range in the color scale has thus been measured. When three colors widely enough separated to form the corners of a color triangle have been reached, such a triangle may be constructed in the usual way from the equation giving their mixture for gray, taking into account also their power of saturation, as already determined. This forms the basis of the system and other colors are assigned places in it in the usual way. For the details of the method and the discussion of the nature of the system thus constructed, the reader is referred to the original. The author unfortunately does not specify how the *hues* to be placed at the corners of this color triangle are to be selected, whether by reference to the spectrum, to well-known pigments, or in some other way—an omission that would have to be supplied before any-